

## Hippuristerones E–I, New Polyoxygenated Steroids from the Gorgonian Coral *Isis hippuris*

Jyh-Horng Sheu,<sup>\*,†</sup> Long-Fei Huang,<sup>†</sup> Shin-Pin Chen,<sup>†</sup> Yi-Lea Yang,<sup>†</sup> Ping-Jyun Sung,<sup>†</sup> Guey-Horng Wang,<sup>†,‡</sup> Jui-Hsin Su,<sup>†</sup> Chih-Hua Chao,<sup>†</sup> Wan-Ping Hu,<sup>§</sup> and Jeh-Jeng Wang<sup>§</sup>

Department of Marine Resources, National Sun Yat-Sen University, Kaohsiung 804, Taiwan, Republic of China, Center for General Education, Hsing-Kuo University, Tainan 709, Taiwan, Republic of China, and School of Chemistry, Kaohsiung Medical University, Kaohsiung 807, Taiwan, Republic of China

Received December 31, 2002


Five novel (22*R*,23*S*,24*S*)-steroids, hippuristerones E–I (**1–5**), have been isolated from the gorgonian coral *Isis hippuris*. The structures of steroids **1–5** were deduced by extensive 1D and 2D NMR studies (<sup>1</sup>H, <sup>13</sup>C, <sup>1</sup>H–<sup>1</sup>H COSY, HMQC, HMBC, and NOESY experiments). The structure of **1** was further supported by molecular mechanics calculations.

Gorgonian corals have been known to be rich sources of polyoxygenated steroids with novel structures that exhibit a variety of biological activities.<sup>1</sup> As part of our ongoing study on chemical constituents of the Taiwanese gorgonian and soft corals,<sup>2</sup> the gorgonian coral *Isis hippuris* L. (phylum Cnidaria, order Gorgonacea, family Isididae),<sup>3</sup> which is a major inhabitant in Taiwanese tropical waters, has been the subject of an investigation. Previous studies on *I. hippuris* have resulted in the isolation of a series of novel metabolites including several highly oxygenated spiroketal steroids<sup>4–8</sup> and polyoxygenated gorgosteroids,<sup>9,10</sup> a (22*R*,23*S*,24*S*)-polyoxygenated steroid, hippuristerone A,<sup>11</sup> and six suberosane-type cytotoxic sesquiterpenes.<sup>12</sup> A study on an Indonesian gorgonian *I. hippuris* also has led to the discovery of a series of polyoxygenated steroids, including hippuristerones B–D.<sup>13</sup> Our present study on the constituents of this gorgonian coral has resulted in the isolation of five novel (22*R*,23*S*,24*S*)-steroids, hippuristerones E–I (**1–5**). The structures, including the relative configurations of the new metabolites **1–5**, were elucidated by extensive spectral analysis, and the structure of **1** was further confirmed by molecular dynamics calculations.

### Results and Discussion

The gorgonian coral *I. hippuris* was frozen immediately after collection and subsequently freeze-dried. The freeze-dried organism was extracted successively with *n*-hexane and CH<sub>2</sub>Cl<sub>2</sub> to afford a crude extract. The crude extract was purified by extensive column chromatography on silica gel and afforded the new steroids **1–5**.

Hippuristerone E (**1**) was obtained as a white powder. The HRFABMS of **1** established a molecular formula of C<sub>33</sub>H<sub>52</sub>O<sub>7</sub>, implying eight degrees of unsaturation. The IR spectrum of **1** showed the presence of hydroxy ( $\nu_{\max}$  3380 cm<sup>-1</sup>) and carbonyl ( $\nu_{\max}$  1720, 1717 cm<sup>-1</sup>) groups in the molecule of **1**. Its <sup>13</sup>C NMR spectrum showed signals of nine methyl, eight methylene, eight methine, and eight quaternary carbons, including those of two ketones ( $\delta$  218.8, s; 211.8, s), two ester carbonyls ( $\delta$  171.4, s; 170.3, s), and three oxygenated sp<sup>3</sup> carbons ( $\delta$  85.7, s; 81.1, d; 77.1, s) (Table 2). By comparison of the above data with those of hippuris-



terone A (**6**),<sup>11</sup> it was found that **1** is a steroid with a structure related to that of **6**. In the <sup>1</sup>H NMR of **1** (Table 1), the doublets at  $\delta$  0.85 (3H, d,  $J$  = 7.0 Hz) and 0.88 ppm (3H, d,  $J$  = 7.0 Hz) were attributed to H<sub>3</sub>-28 and H<sub>3</sub>-29. Furthermore, five singlets appearing at  $\delta$  1.45 (3H), 1.43 (3H), 1.27 (3H), 1.04 (3H), and 1.00 ppm (3H) were due to the resonances of H<sub>3</sub>-26, H<sub>3</sub>-27, H<sub>3</sub>-21, H<sub>3</sub>-19, and H<sub>3</sub>-18, respectively. Two signals that appeared at  $\delta$  2.12 (3H, s) and d 1.96 ppm (3H, s) revealed the presence of two acetoxy groups. From the <sup>1</sup>H–<sup>1</sup>H COSY spectrum of **1** (Figure 1), it was possible to establish the proton sequences from H<sub>2</sub>-1 to H<sub>2</sub>-2; H<sub>2</sub>-4 to H-9; H-8 to H-14; H-9 to H<sub>2</sub>-11; H<sub>2</sub>-11 to H<sub>2</sub>-12; H-14 to H<sub>2</sub>-15; H-22 to H-24; H-23 to H<sub>3</sub>-29; and H-24 to H<sub>3</sub>-28. The ring-junctured C-18 and C-19 methyl groups were positioned at C-10 and C-13, respectively, from the key HMBC correlations of H<sub>3</sub>-19 with C-1, C-5, C-9, and C-10, and H<sub>3</sub>-18 with C-12, C-13, C-14, and C-17,

\* To whom correspondence should be addressed. Tel: 886-7-5252000, ext. 5030. Fax: 886-7-5255020. E-mail: sheu@mail.nsysu.edu.tw.

<sup>†</sup> National Sun Yat-Sen University.


<sup>‡</sup> Hsing-Kuo University.

<sup>§</sup> Kaohsiung Medical University.

**Table 1.**  $^1\text{H}$  NMR Chemical Shifts for Steroids **1**–**5**

proton	<b>1</b> <sup>a</sup>	<b>2</b> <sup>b</sup>	<b>3</b> <sup>c</sup>	<b>4</b> <sup>c</sup>	<b>5</b> <sup>c</sup>
1	2.02 m; 1.33 m	1.99 m; 1.34 m	2.00 m; 1.40 m	2.05 m; 1.33 m	2.04 m; 1.62 m
2	2.40 m; 2.22 m	2.30 m; 2.38 m	2.38 m	2.30 m	2.32 m
4	2.28 m; 2.11 m	2.27 m; 2.11 m	2.30 m; 2.10 m	2.27 m; 2.10 m	2.27 m; 2.09 m
5	1.54 m	1.56 m	1.56 m	1.56 m	1.54 m
6	1.38 m	1.40 m	1.40 m	1.40 m	1.37 m
7	1.65 m; 1.02 m	1.81 m; 0.91 m	1.80 m; 0.92 m	1.82 m; 0.92 m	1.78 m; 0.92 m
8	1.60 m	1.71 m	1.76 m	1.56 m	1.49 m
9	0.93 m	0.78 m	0.89 m	0.70 m	0.75 m
11	1.60 m; 1.48 m	1.65 m	1.52 m; 1.34 m	1.60 m; 1.48 m	1.59 m; 1.37 m
12	2.26 m; 1.49 m	2.17 m; 1.37 m	2.15 m; 1.40 m	1.82 m; 1.48 m	1.88 m; 1.46 m
14	1.45 m	1.22 m	1.42 m	1.06 m	1.29 m
15	2.28 m; 1.90 m	2.22 m	2.28 m; 1.45 m	2.20 m; 1.45 m	1.64 m; 1.40 m
16		4.09 t (7.0)	4.05 t (6.9)	4.06 t (7.2)	2.08 m; 1.91 m
17	2.20 s				
18	1.00 s	4.31 d (11.5); 4.22 d (11.5)	4.31 d (11.4); 4.21 d (11.4)	0.95 s	0.92 s
19	1.04 s	1.03 s	1.02 s	1.01 s	1.00 s
21	1.27 s	1.61 s	1.59 s	1.59 s	1.55 s
22	5.36 d (2.8) <sup>d</sup>	4.64 d (10.5)	4.67 d (10.7)	4.62 d (10.8)	4.73 d (10.7)
23	2.22 m	1.48 m	2.30 m	2.45 m	2.33 m
24	1.92 m	1.54 m	2.05 m	1.55 m	1.59 m
26	1.45 s	1.24 s	1.56 s	1.23 s	1.28 s
27	1.43 s	1.21 s	1.42 s	1.20 s	1.20 s
28	0.85 d (7.0)	0.90 d (7.0)	0.90 d (7.4)	0.88 d (7.2)	0.91 d (7.4)
29	0.88 d (7.0)	0.86 d (7.0)	0.87 d (6.6)	0.84 d (6.9)	0.82 d (6.6)
OH-16				3.27 s	
OH-20	3.49 br s				
acetate	2.12 s;	2.13 s;	2.12 s;	2.11 s;	2.08 s;
methyls	1.96 s	2.07 s	2.07 s; 1.98 s		

<sup>a</sup> Spectra recorded at 400 MHz in  $\text{CDCl}_3$  at 25 °C. <sup>b</sup> Spectra recorded at 500 MHz in  $\text{CDCl}_3$  at 25 °C. <sup>c</sup> Spectra recorded at 300 MHz in  $\text{CDCl}_3$  at 25 °C. <sup>d</sup>  $J$  values (in Hz) in parentheses. The values are ppm downfield from TMS.

**Figure 1.** Selective HMBC and  $^1\text{H}$ – $^1\text{H}$  COSY correlations of **1**.

observed in an HMBC experiment (Figure 1). The  $^{13}\text{C}$  NMR signals at  $\delta$  171.4 (s) and 170.3 ppm (s) correlated with the signals of the methyl protons at  $\delta$  2.12 (3H, s) and 1.96 ppm (3H, s) in the HMBC spectrum of **1** (Figure 1) and were consequently assigned as the carbon atoms of the two acetate carbonyls. The HMBC experiment of **1** further revealed the connectivity between H-22 ( $\delta$  5.36, 1H, d,  $J$  = 2.8 Hz) and the carbonyl carbon ( $\delta$  171.4, s) of an acetate unit and demonstrated the location of an acetoxy group to be at C-22. On the basis of the consideration of the molecular formula and by comparison of the NMR spectral data with **6**, the second acetoxy group should be attached at C-25 and one hydroxy group had to be placed at C-20. The two ketone groups at C-3 and C-16 were confirmed by their HMBC correlations with H<sub>2</sub>-2 and H<sub>2</sub>-4, and H<sub>2</sub>-15

and H-17, respectively. The above observations and several other  $^1\text{H}$ – $^1\text{H}$  COSY and HMBC correlations (Table 3 and Figure 1) thus provided unambiguous evidence for the molecular framework of **1**.

The relative stereochemistry of **1** was deduced using a NOESY experiment (Table 3). In the NOESY spectrum of **1**, H<sub>3</sub>-21 did not give correlation with H<sub>3</sub>-18, and H-17 was found to exhibit correlations with H<sub>3</sub>-21 and H-14. Thus, H-14, H-17, and H<sub>3</sub>-21 should be placed on the  $\alpha$ -face, since the C-18 methyl is the  $\beta$ -substituent at C-13, and the hydroxy group at C-20 should be  $\beta$ -oriented. Furthermore, H-8 exhibited NOE correlations with H<sub>3</sub>-18 and H<sub>3</sub>-19, but not with H-9 and H-14, indicating that H<sub>3</sub>-19 and H-8 are situated on the  $\beta$ -face, and H-9 is situated on the  $\alpha$ -face. Also, H-5 was found to exhibit correlations with H-9, but not with H<sub>3</sub>-19, indicating that H-5 was  $\alpha$ -oriented in **1**. The relative stereochemistry of the side chain of **1** is also determined by a NOESY experiment (Table 3). H-17 was found to show NOE correlations with H-22, H-23, and H-24. By detailed consideration of molecular models, it was suggested that the side chain substituents of **1** should possess orientations of C-22 $\alpha$ , C-23 $\alpha$ , and C-24 $\alpha$ . Thus, hippuristerone E (**1**) is considered to be a novel 20-hydroxysteroid possessing a (22*R*,23*S*,24*S*)-23,24-dimethyl-22,25-diacetoxy side chain subunit.


Geometry optimization was performed using DISCOVER utilizing the CVFF (consistent valence force field) calculations for energy minimization. The results were visualized using INSIGHT II running on a Silicon Graphics IRIS

**Table 2.**  $^{13}\text{C}$  NMR Chemical Shifts for Steroids **1**–**5**

carbon	<b>1</b> <sup>a</sup>	<b>2</b> <sup>b</sup>	<b>3</b> <sup>c</sup>	<b>4</b> <sup>c</sup>	<b>5</b> <sup>c</sup>
1	38.1 t	38.3 t	38.4 t	38.4 t	38.5 t
2	38.0 t	38.0 t	38.1 t	38.1 t	38.2 t
3	211.8 s	211.6 s	211.6 s	211.8 s	211.9 s
4	44.5 t	44.5 t	44.6 t	44.6 t	44.7 t
5	46.5 d	46.5 d	46.6 d	46.6 d	46.6 d
6	28.6 t	28.5 t	28.7 t	28.7 t	28.8 t
7	31.5 t	31.6 t	31.7 t	31.6 t	31.5 t
8	33.8 d	34.4 d	34.5 d	34.7 d	35.6 d
9	53.4 d	53.8 d	53.3 d	53.7 d	53.6 d
10	35.7 s	35.7 s	35.7 s	35.7 s	35.7 s
11	20.9 t	21.5 t	21.6 t	21.4 t	21.6 t
12	39.3 t	32.4 t	33.5 t	36.7 t	36.5 t
13	43.6 s	45.6 s	45.6 s	43.1 s	43.8 s
14	50.9 d	49.3 d	48.2 d	49.3 d	54.9 d
15	38.8 t	33.4 t	32.0 t	33.4 t	23.6 t
16	218.8 s	70.1 d	70.2 d	70.2 d	30.9 t
17	67.4 d	77.3 s	77.8 s	79.2 s	79.1 s
18	14.4 q	63.4 t	63.5 t	15.6 q	15.4 t
19	11.4 q	11.4 q	11.5 q	11.4 q	11.5 q
20	77.1 s	66.8 s	66.5 s	67.8 s	67.3 s
21	20.6 q	16.2 q	16.3 q	16.5 q	17.0 q
22	81.1 d	77.3 d	77.2 d	77.7 d	78.5 d
23	32.3 d	33.1 d	33.8 d	33.1 d	33.4 d
24	44.5 d	41.8 d	40.0 d	41.8 d	41.7 d
25	85.7 s	73.8 s	85.7 s	73.8 s	73.8 s
26	23.6 q	30.8 q	23.4 q	30.8 q	30.5 q
27	25.1 q	25.9 q	25.2 q	26.1 q	26.3 q
28	9.5 q	11.4 q	10.5 q	11.5 q	11.3 q
29	10.7 q	12.0 q	12.0 q	12.1 q	11.9 q
acetate methyls	22.7 q	21.2 q	21.1 q	21.1 q	21.1 q
			21.2 q		
			22.7 q		
acetate carbonyls	171.4 s	171.4 s	171.3 s	171.6 s	170.7 s
	170.3 s	171.3 s	171.1 s	171.1 s	171.1 s
			169.9 s	169.9 s	169.9 s

<sup>a</sup> Spectra recorded at 100 MHz in  $\text{CDCl}_3$  at 25 °C. <sup>b</sup> Spectra recorded at 125 MHz in  $\text{CDCl}_3$  at 25 °C. <sup>c</sup> Spectra recorded at 75 MHz in  $\text{CDCl}_3$  at 25 °C. <sup>d</sup> Multiplicity deduced by DEPT and indicated by usual symbols. The values are ppm downfield from TMS.

(SGI) INDIGO XS24-4000. The conformational search suggested the most stable conformation as shown in Figure 2. The most stable conformation revealed the presence of a hydrogen bond between the hydroxy group at C-20 and the oxygen at C-22 (the H–O distance is 2.43 Å), so that the conformation around C-20/C-22 bond is fixed, and in turn prevents the free rotation around the C-22/C-23 bond. The distance between H-17 and H-24 was found to be 4.09 Å, which could reasonably explain the NOE correlations observed between these two protons. Furthermore, the orthogonal (torsional angle around 90°) arrangement of H-22 and H-23 protons can be used to rationalize the

**Figure 2.** Stereoview of **1** generated from computer modeling.


presence of a very small coupling constant (2.8 Hz) between these two methine protons in the  $^1\text{H}$  NMR spectrum of **1** (Table 1). On the basis of the above results, the molecular structure of **1** was established unambiguously.

Hippuristerone F (**2**) was obtained as a white amorphous solid that gave a  $[\text{M} + \text{H}]^+$  peak at  $m/z$  577.3742 in the HRFABMS, appropriate for a molecular formula of  $\text{C}_{33}\text{H}_{52}\text{O}_8$ , requiring eight degrees of unsaturation. The LRFABMS showed peaks at  $m/z$  559  $[\text{M} + \text{H} - \text{H}_2\text{O}]^+$ , 541  $[\text{M} + \text{H} - 2 \text{H}_2\text{O}]^+$ , 517  $[\text{M} + \text{H} - \text{AcOH}]^+$ , 457  $[\text{M} + \text{H} - 2 \text{AcO}]^+$ , and 421  $[\text{M} + \text{H} - 2 \text{AcOH} - 2 \text{H}_2\text{O}]^+$ , suggesting the presence of two hydroxy and two acetoxy groups in the molecular structure of **2**. The  $^{13}\text{C}$  NMR spectrum of **2** showed signals of eight methyl, nine methylene, eight methine, and eight quaternary carbons, including one ketone ( $\delta$  211.6, s), two ester carbonyls ( $\delta$  171.4, s; 171.3, s), six carbons bonded to an oxygen ( $\delta$  77.3, s; 77.3, d; 73.8, s; 70.1, d; 66.8, s; 63.4, t), and two normal quaternary carbons ( $\delta$  45.6; 35.7) (Table 2). It was found that the above data are very similar to the carbon shifts of hippuristerone A (**6**), indicating that the structure of **2** should be close to **6**. However, it was observed that the signal of the methyl carbon C-18 in molecules of **2** disappeared and was replaced by a signal resonating at  $\delta$  63.4 (t). Also, the resonance of H<sub>3</sub>-18 in **6** was replaced by signals that were downfield shifted to  $\delta$  4.31 (1H, d,  $J = 11.5$  Hz) and 4.22 (1H, d,  $J = 11.5$  Hz). Thus, the methyl group attached at C-13 in **6** should be converted to an oxygen-bearing methylene group. The  $^{13}\text{C}$  NMR signals resonating at 171.4 and 171.3 ppm were found to be correlated with the signals of the methyl protons at  $\delta$  2.13 (3H, s) and 2.07 (3H, s) in the HMBC spectrum of **2** and consequently assigned as the carbon atoms of the two acetate carbonyls. The HMBC spectrum of **2** further revealed the connectivities between H-22 ( $\delta$  4.64, 1H, d,  $J = 10.5$  Hz) and a carbonyl carbon ( $\delta$  171.4),

**Table 3.** Selective  $^1\text{H}$ – $^1\text{H}$  COSY, HMBC, and NOESY Correlations for **1**

C/H	$^1\text{H}$ – $^1\text{H}$ COSY	HMBC	NOESY
3		H <sub>2</sub> -2, H <sub>2</sub> -4	
16		H <sub>2</sub> -15, H-17	
17		H <sub>3</sub> -18, H <sub>3</sub> -21	
18		H <sub>2</sub> -12, H-14, H-17	H-14, H <sub>3</sub> -21, H-22, H-23, H-24
19		H <sub>2</sub> -2, H-5, H-9	H-8
20		H-17, H <sub>3</sub> -21, H-22	H-8
21		H-17, H-22	H-17, H-22
22	H-23	H-17, H <sub>3</sub> -21, H-23, H-24, H <sub>3</sub> -29	H-17, H <sub>3</sub> -21, H-23, H-24
23	H-22, H-24, H <sub>3</sub> -29	H-22, H-24, H <sub>3</sub> -28, H <sub>3</sub> -29	H-17, H-22
24	H-23, H <sub>3</sub> -28	H-22, H-23, H <sub>3</sub> -26, H <sub>3</sub> -27, H <sub>3</sub> -28, H <sub>3</sub> -29	H-17, H-22
25		H-24, H <sub>3</sub> -26, H <sub>3</sub> -27, H <sub>3</sub> -28	
26		H-24	
27		H-24	
28	H-24	H-23, H-24	
29	H-23	H-22, H-23, H-24	
22-OCOMe		H-22	





**Figure 3.** Possible biosynthetic pathway of hippuristerone (**1**).

and between H<sub>2</sub>-18 and the other carbonyl carbon ( $\delta$  171.3), demonstrating the locations of two acetoxy groups to be at C-22 and C-18. On the basis of these results, the structure of hippuristerone F was established as 25-deacetoxy-25-hydroxy-18-acetoxyhippuristerone A, as described by formula **2**.

Hippuristerone G (**3**) was obtained as a white powder. The molecular formula C<sub>35</sub>H<sub>54</sub>O<sub>9</sub> was established by a FAB mass spectrum, which gave a [M + H]<sup>+</sup> peak at *m/z* 619, and by <sup>13</sup>C NMR spectral data, which revealed the presence nine methyl, nine methylene, eight methine, and nine quaternary carbons, including one ketone ( $\delta$  211.6, s), three ester carbonyls ( $\delta$  171.3, s; 171.1, s; 169.9, s), six carbons attached by an oxygen ( $\delta$  85.7, s; 77.8, s; 77.2, d; 70.2, d; 66.5, s; 63.5, t), and two normal quaternary carbons ( $\delta$  45.6; 35.7). In comparison of the <sup>13</sup>C and <sup>1</sup>H NMR spectral data of **3** with those of **2** and **6**, the structure of hippuristerone G was established as 18-acetoxyhippuristerone A, as described by formula **3**.

Hippuristerones H (**4**) and I (**5**) had molecular formulas of C<sub>31</sub>H<sub>50</sub>O<sub>6</sub> and C<sub>31</sub>H<sub>50</sub>O<sub>5</sub>, respectively, as suggested by their NMR (Tables 1 and 2) and HRFASMS data. In comparison of both <sup>1</sup>H and <sup>13</sup>C NMR spectral data of these two metabolites with those of hippuristerone A (**6**), it was concluded that the structures of **4** and **5** are very similar to that of **6**, except that the acetoxy group attached at C-25 of **6** was replaced by a hydroxy group in both **4** and **5**. Also, it was observed that the hydroxymethine functionality at C-16 of **4** was reduced to the corresponding methylene moiety in **5**. Thus, the structures of **4** and **5** were established unambiguously as 25-deacetoxy-25-hydroxyhippuristerone A and 16-dehydroxy-25-deacetoxy-25-hydroxyhippuristerone A, respectively.

Hippuristerone E (**1**) appeared to be biosynthesized from hippuristerone A (**6**) by a pathway as shown in Figure 3. The 17 $\beta$ ,20 $\beta$ -epoxy group of **6** was ring-opened to form the intermediate **7**, which has a carbonium ion at C-17 and a  $\beta$ -hydroxy group at C-20. The Wagner–Meerwein shift of H $\alpha$ -16 to the  $\alpha$ -face of C-20 and the following deprotonation of the 16-hydroxy group converted the intermediate **7** into **1**.

## Experimental Section

**General Experimental Procedures.** Melting points were determined using a Fisher-Johns melting point apparatus and have not been corrected. Optical rotations were measured on a Jasco DIP-1000 digital polarimeter. IR spectra were recorded on Hitachi I-2001 and Jasco FT/IR-5300 infrared spectrophotometers. The NMR spectra were recorded on FT-NMR instruments at 300, 400, or 500 MHz for <sup>1</sup>H and 75, 100, or 125 MHz for <sup>13</sup>C, respectively, in CDCl<sub>3</sub> using TMS as an internal standard. Nuclear Overhauser and exchange spectroscopy (NOESY), <sup>1</sup>H–<sup>1</sup>H correlation spectroscopy (COSY), heteronuclear multiple quantum coherence (HMQC), and <sup>1</sup>H-detected multiple-bond heteronuclear multiple quantum coherence (HMBC) experiments were performed by using standard Bruker pulse sequences. Low-resolution mass spectra

were obtained by fast atom bombardment (FAB) on a VG Quattro GC/MS spectrometer. High-resolution mass spectra (HRMS) were recorded by fast atom bombardment on a JEOL JMX-HX 110 mass spectrometer. Silica gel 60 (Merck, 230–400 mesh) was used for column chromatography. Precoated silica gel plates (Merck Kieselgel 60 F<sub>254</sub>, 2 mm) were used for analytical TLC.

**Animal Material.** The gorgonian coral *I. hippuris* was collected by hand using scuba at Green Island, which is located off the southeast coast of Taiwan, in February 1999, at a depth of 25 m, and was stored in a freezer until extraction. A voucher specimen was deposited in the Department of Marine Resources, National Sun Yat-Sen University (specimen no. GISC-102).

**Extraction and Isolation.** The gorgonian coral (4.3 kg fresh wt) was collected and freeze-dried. The freeze-dried material was minced and extracted exhaustively with *n*-hexane and CH<sub>2</sub>Cl<sub>2</sub>. The combined organic extract was evaporated to give a dark green residue (37.0 g), which was chromatographed on a SiO<sub>2</sub> column using solvents of increasing polarity from *n*-hexane to EtOAc. Steroids **1** (12.9 mg) and **2** (2.7 mg) were eluted with *n*-hexane/EtOAc (3:1). Steroid **3** (6.3 mg) was obtained by elution with *n*-hexane/EtOAc (5:1). A fraction eluted with *n*-hexane/EtOAc (3:1–3:2) was further purified by SiO<sub>2</sub> column using acetone/CH<sub>2</sub>Cl<sub>2</sub> (1:3) to afford **4** (7.9 mg). A fraction eluted with *n*-hexane/EtOAc (4:1–3:1) was further purified by SiO<sub>2</sub> column using acetone/CH<sub>2</sub>Cl<sub>2</sub> (1:3) to afford **5** (14.5 mg).

**Hippuristerone E (1):** white powder; mp 174–176 °C; [ $\alpha$ ]<sub>D</sub> –92° (c 0.1, CHCl<sub>3</sub>); IR (KBr)  $\nu_{\max}$  3380, 1720, and 1717 cm<sup>-1</sup>; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Tables 1 and 2; FABMS *m/z* 583 [(M + Na)<sup>+</sup>, 2], 565 (0.2), 523 (1), 483 (1), 439 (1), 423 (1), 391 (3), and 341 (7); HRFABMS *m/z* 583.3626 (M + Na)<sup>+</sup> (calcd for C<sub>33</sub>H<sub>52</sub>O<sub>7</sub>Na, 583.3611).

**Hippuristerone F (2):** white powder; mp 125–127 °C; [ $\alpha$ ]<sub>D</sub> –15° (c 0.3, CHCl<sub>3</sub>); IR (KBr)  $\nu_{\max}$  3452, 1726, and 1244 cm<sup>-1</sup>; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Tables 1 and 2; FABMS *m/z* 577 [(M + H)<sup>+</sup>]; HRFABMS *m/z* 577.3742 (M + H)<sup>+</sup> (calcd for C<sub>33</sub>H<sub>53</sub>O<sub>8</sub>, 577.3726).

**Hippuristerone G (3):** white powder; mp 113–114 °C; [ $\alpha$ ]<sub>D</sub> 23° (c 0.02, CHCl<sub>3</sub>); IR (KBr)  $\nu_{\max}$  3495, 1728, and 1248 cm<sup>-1</sup>; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Tables 1 and 2; FABMS *m/z* 619 [(M + H)<sup>+</sup>, 0.01], 551 (0.04), 499 (0.01), 481 (0.01), 439 (0.02), and 421 (0.02); HRFABMS *m/z* 641.3666 (M + Na)<sup>+</sup> (calcd for C<sub>35</sub>H<sub>54</sub>O<sub>9</sub>Na, 641.3667).

**Hippuristerone H (4):** white powder; mp 122–124 °C; [ $\alpha$ ]<sub>D</sub> 9° (c 0.31, CHCl<sub>3</sub>); IR (KBr)  $\nu_{\max}$  3493, 1726, and 1711 cm<sup>-1</sup>; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Tables 1 and 2; FABMS *m/z* 541 [(M + Na)<sup>+</sup>]; HRFABMS *m/z* 519.3682 (M + H)<sup>+</sup> (calcd for C<sub>31</sub>H<sub>51</sub>O<sub>6</sub>, 519.3672).

**Hippuristerone I (5):** white powder; mp 120–122 °C; [ $\alpha$ ]<sub>D</sub> 12° (c 0.36, CHCl<sub>3</sub>); IR (KBr)  $\nu_{\max}$  1730, 1709, and 1246 cm<sup>-1</sup>; <sup>1</sup>H NMR and <sup>13</sup>C NMR data, see Tables 1 and 2; FABMS *m/z* 525 [(M + Na)<sup>+</sup>]; HRFABMS *m/z* 503.3734 (M + H)<sup>+</sup> (calcd for C<sub>31</sub>H<sub>51</sub>O<sub>5</sub>, 503.3723).

**Molecular Mechanics Calculations.** The minimum energy conformation of hippuristerone E (**1**) was determined using the MSI Insight II/DISCOVER version 95 molecular modeling package incorporating an empirical force field, the consistent valence force field (CVFF),<sup>14</sup> on a Silicon Graphics IRIS Indigo XS24/R4000 workstation. All the force field calculations were carried out in vacuo (dielectric constant = 1). The conformational space of steroid **1** was scanned using the high-temperature molecular dynamics simulation technique followed by energy minimization. A 100 ps molecular dynamics simulation at 1000 K provided a set of 500 conformations of **1**. Each of them was used as a starting structure for the subsequent energy minimization (1000 steps, conjugated gradient algorithm). In the subsequent analysis, only the 15 conformations with a reasonably low energy (at most 5 kcal/mol higher with respect to the lowest energy conformer) were used. The conformer shown in Figure 2 is the lowest energy conformation of steroid **1**.

**Acknowledgment.** This work was supported by a grant from the National Science Council of the Republic of China (Contract No. NSC-89-2113-M-110-025) awarded to J.-H.S.

### References and Notes

- (1) Faulkner, D. J. *Nat. Prod. Rep.* **2001**, *18*, 1–49, and previous reports in this series.
- (2) Wang, G.-H.; Sheu, J.-H.; Duh, C.-Y.; Chiang, M. Y. *J. Nat. Prod.* **2002**, *65*, 1475–1478, and references therein.
- (3) Bayer, F. M. *Proc. Biol. Soc. Wash.* **1981**, *94*, 902–947.
- (4) Kazlauskas, R.; Murphy, P. T.; Quinn, R. J.; Wells, R. J.; Schönholzer, P. *Tetrahedron Lett.* **1977**, 4439–4442.
- (5) Higa, T.; Tanaka, J.; Tsukitani, Y.; Kikuchi, H. *Chem. Lett.* **1981**, 1647–1650.
- (6) Higa, T.; Tanaka, J.; Tachibana, K. *Tetrahedron Lett.* **1981**, *22*, 2777–2780.
- (7) Rao, C. B.; Ramana, K. V.; Rao, D. V.; Fahy, E.; Faulkner, D. J. *J. Nat. Prod.* **1988**, *51*, 954–958.
- (8) Tanaka, J.; Trianto, A.; Musman, M.; Issa, H. H.; Ohtani, I. I.; Ichiba, T.; Higa, T.; Yoshida, W. Y.; Scheuer, P. J. *Tetrahedron* **2002**, *58*, 6259–6266.
- (9) Tanaka, J.; Higa, T.; Tachibana, K.; Iwashita, T. *Chem. Lett.* **1982**, 1295–1296.
- (10) Shen, Y.-C.; Prakash, C. V. S.; Chang, Y.-T. *Steroids* **2001**, *66*, 721–725.
- (11) Sheu, J.-H.; Chen, S.-P.; Sung, P.-J.; Chiang, M. Y.; Dai, C.-F. *Tetrahedron Lett.* **2000**, *41*, 7885–7888.
- (12) Sheu, J.-H.; Hung, K.-C.; Wang, G.-H.; Duh, C.-Y. *J. Nat. Prod.* **2000**, *63*, 1603–1607.
- (13) González, N.; Barral, M. A.; Rodríguez, J.; Jiménez, C. *Tetrahedron* **2001**, *57*, 3487–3497.
- (14) MSI Insight II/DISCOVER (version 95.0/2.97) is a molecular modeling software package of MSI Technologies Inc., Barnes Canyon Rd., San Diego, CA 92121.

NP020602R