

## Original Article

## Excoecarianin, Isolated from *Phyllanthus urinaria* Linnea, Inhibits Herpes Simplex Virus Type 2 Infection through Inactivation of Viral Particles

Hua-Yew Cheng<sup>1</sup>, Chien-Min Yang<sup>2,3</sup>, Ta-Chen Lin<sup>4</sup>, Liang-Tzung Lin<sup>5</sup>, Lien-Chai Chiang<sup>6</sup> and Chun-Ching Lin<sup>2</sup>

<sup>1</sup>Department of Cosmetic Applications & Management, Tung Fang Institute of Technology, 829 Kaohsiung County, <sup>2</sup>School of Pharmacy, College of Pharmacy, Kaohsiung Medical University, 807 Kaohsiung, <sup>3</sup>Department of Cosmetology and Health Care, Min Hwei College of Health Care Management, 736 Tainan County, <sup>4</sup>Department of Nursing, Central Taiwan University of Sciences and Technology, 406 Taichung, Taiwan, <sup>5</sup>Department of Microbiology & Immunology, Dalhousie University, Halifax, Nova Scotia, Canada B3H 1X5 and <sup>6</sup>Department of Microbiology, College of Medicine, Kaohsiung Medical University, 807 Kaohsiung, Taiwan

*Phyllanthus urinaria* Linnea (Euphorbiaceae) is one of the traditional medicinal plants widely used by oriental people to treat various diseases. We have previously demonstrated that the acetone extract of *P. urinaria* inhibits herpes simplex virus type 2 (HSV-2) but not HSV-1 infection. In a continuing effort to clarify the antiviral mechanisms of *P. urinaria*, we isolated the pure compound excoecarianin from the whole plant of *P. urinaria* through acetone extraction, and investigated its anti-HSV-1 and HSV-2 activities. Our results indicated that excoecarianin protected Vero cells from HSV-2 but not HSV-1 infection, and its 50% inhibitory concentration (IC<sub>50</sub>) was 1.4 ± 0.1 μM. The antiviral effective concentration of excoecarianin did not affect the viability or the morphology of Vero cells. Although excoecarianin inhibited HSV-2 infection, the inhibitory effect, however, was most prominent when excoecarianin was concurrently added with the virus. Pretreatment of Vero cells with excoecarianin with removal of the drug prior to infection did not yield any antiviral effects, and the same observation was made for post viral entry treatment. Subsequent studies revealed that excoecarianin inactivated HSV-2 virus particles to prevent viral infection. A synergistic antiviral effect against HSV-2 was also observed when Vero cells were treated with a combination of acyclovir (ACV) and excoecarianin. These results suggested that excoecarianin merits to be further explored as an entry inhibitor against HSV-2 and could potentially be investigated for combinatorial drug treatment with nucleoside analogues such as ACV in therapeutic management of HSV-2 infection.

**Keywords:** antiviral activity – excoecarianin – herpes simplex virus type 2 – *Phyllanthus urinaria* – viral inactivation

### Introduction

Herpes simplex virus type 1 (HSV-1) and type 2 (HSV-2) are two species of the *Herpesviridae* family, which

consists of large DNA-enveloped viruses. While HSV-1 is more frequently associated with oral–facial mucocutaneous lesions (cold sores and fever blisters), HSV-2 is more commonly associated with genital herpes. The two viruses can cause a variety of diseases, which in certain cases, may lead to complications, especially in immunocompromised patients. Herpes viruses usually produce

For reprints and all correspondence: C.-C. Lin, No. 100, Shin-Chuan 1st Road, Kaohsiung Medical University, Kaohsiung City, Taiwan.  
Tel: +886-7-3121101 ext. 2122; Fax: +886-7-3135125;  
E-mail: aalin@kmu.edu.tw

© The Author(s) 2009. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/2.5/uk/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

lifelong infections, and recurrent infection may occur upon viral reactivation through stimuli such as sunlight, stress and weakened immunity.

Nucleoside analogs such as acyclovir (ACV) are the primary candidates used to treat HSV infections. Although some non-nucleoside inhibitors of herpesviruses have been developed (1–3), few of them are officially approved for HSV therapy (4,5). The lack of approved therapeutics and the emergence of ACV-resistant HSV strains have made the management of HSV infection a challenge, particularly in immunocompromised patients. In fact, previous reports have indicated that ACV-resistant HSV is more frequently identified in immunocompromised individuals rather than in those who are immunocompetent (6,7). Consequently, there is still a need to search for novel and more effective antiviral agents to prevent and/or to treat HSV infections (8–10).

Many naturally occurring plants, either as extracts or as pure compounds, have been reported to exhibit antiviral activities (11–14), indicating that plant materials may serve as a source for the discovery of antiviral agents. *Phyllanthus urinaria* Linnaea (Euphorbiaceae) is a traditional medicinal plant that is widely used in Eastern Asia to treat a variety of diseases. The pharmacological activities of *P. urinaria* have been well documented in literature (15–36), and include antiviral activities (37–43) as well.

In our previous studies, we showed that the acetone, ethanol and methanol extracts of *P. urinaria* inhibited HSV infections *in vitro* (44). Further isolation of gemin D, geraniin, hippomanin A and 1, 3, 4, 6-tetra-*O*-galloyl- $\beta$ -D-glucose from the acetone extract of *P. urinaria* was found to suppress HSV infections at different magnitudes of potency (45,46). In a continuing effort to clarify the antiviral profile of *P. urinaria*, we isolated another pure compound, excoecarianin, from the whole plant of *P. urinaria* by acetone extraction, and investigated its anti-HSV-1 and HSV-2 activities.


## Methods

### Plant Materials

The plant *P. urinaria* was collected from Ping-Tung County and was authenticated using morphological and anatomical techniques. A voucher specimen, with reference number of KMU-HL-PUL 2002, was deposited at the Herbarium of the Graduate Institute of Natural Products of Kaohsiung Medical University.

### Isolation of Excoecarianin from *P. urinaria*

Total 9.5 kg of fresh whole *P. urinaria* plant was cut into small pieces and extracted with acetone–water (4:1, v/v).



**Figure 1.** Chemical structure of excoecarianin.

The extract was concentrated under reduced pressure and then filtered. The filtrate was subsequently eluted with water–methanol and then with water–acetone through Sephadex LH-20 to give four fractions. Fraction 4 was further chromatographed on Sephadex LH-20, MCI-gel CHP 209, Fuji gel ODSG3, Sephadex LH-20 and Bondapak C18/porasil B to get 101 mg of excoecarianin (Fig. 1) with the yield of 0.001%. The structure and purity (>95%) of excoecarianin were determined by spectroscopic and physical data analysis (47).

ACV and excoecarianin were dissolved in dimethyl sulfoxide (DMSO) and then diluted with sterile de-ionized distilled water before use. The final concentration of DMSO was <0.1%, which was not toxic to Vero cells as shown previously (44). A 0.1% DMSO solution was included as control during the experiments.

### Cell and Viruses

African green monkey kidney cells (Vero) (ATCC CCR-81) were used for antiviral assays. Vero cells were

propagated in DMEM (Gibco, Invitrogen, USA) containing 5% of fetal calf serum (FCS). HSV-1 KOS strain and HSV-2 196 strain were grown in Vero cells. The viral titer was determined by plaque assay according to previously described procedures (48) and was expressed as plaque forming units (pfu) per milliliter. Virus stocks were stored at  $-80^{\circ}\text{C}$  until use.

### Cytotoxic Assay

Excoecarianin was tested for its cytotoxic effect on Vero cells by XTT (Sodium 3'-[1-(phenylamino-carbonyl)-3,4-tetrazolium]-bis (4-methoxy-6-nitro) benzene sulfonic acid) (Sigma-Aldrich, USA) assay as described previously (45). Briefly, Vero cells were seeded onto 96-well culture plates (Falcon, BD Biosciences, USA) at  $1 \times 10^4$  cells per well. After 4 h of incubation to allow the cells to settle, excoecarianin was added into each well at various concentrations. The plate was then incubated at  $37^{\circ}\text{C}$  in an atmosphere of 5%  $\text{CO}_2$  for 72 h. Later, the medium was discarded, and the cells were subsequently washed with phosphate buffered saline (PBS). The XTT reagent at a concentration of  $300 \mu\text{g/ml}$  was added, and the plate was re-incubated at  $37^{\circ}\text{C}$  for an additional 2 h to allow the development of formazan. The optical densities (ODs) were then measured with enzyme immunoassay (EIA) reader (Lab Systems MTX Labs, USA) at a test wavelength of 492 nm and a reference wavelength of 690 nm. The cytotoxic effect of excoecarianin on Vero cells was evaluated according to the OD readings obtained and the 50% cytotoxic concentration ( $\text{CC}_{50}$ ) was calculated as previously described (45,46). Besides cytotoxic cell death, Vero cells were also monitored in the presence of excoecarianin for changes in cellular morphology for up to 7 days.

### Antiviral Assay

The inhibitory effect of excoecarianin on HSV infection was investigated by plaque reduction assay (49). Vero cells were seeded onto 24-well culture plates (Falcon, BD Biosciences, USA) at a density of  $1 \times 10^5$  cells per well and incubated for 48 h to reach at least 95% confluency. The medium was then aspirated, and the cell monolayer was infected with 100 pfu of HSV-1 or HSV-2 in the absence or presence of excoecarianin. After 1 h incubation to allow virus adsorption, the cell monolayer was overlaid with overlay medium containing 1% of methylcellulose. The plate was further incubated at  $37^{\circ}\text{C}$  in an atmosphere of 5%  $\text{CO}_2$  for 48 h. Later, the overlay medium was removed, and the infected cell monolayer was fixed with 10% formalin. The cell monolayer was then stained with 1% crystal violet. The fraction of percent inhibition in inhibiting HSV infection was determined, and the minimal concentration that

inhibited the formation of virus plaque number by 50% ( $\text{IC}_{50}$ ) was calculated (49).

### Time-of-addition Study

The time-of-addition effect of excoecarianin was examined according to the previously described procedures with some modifications (49). Briefly, Vero cells were seeded onto 24-well culture plates (Falcon, BD Biosciences, USA) at a density of  $1 \times 10^5$  cells per well and incubated for 48 h to reach at least 95% confluency. Excoecarianin, at concentration of  $1.5 \mu\text{M}$ , was then added onto the cells at either before ( $-6$  and  $-2$  h), during (0 h) or after (2, and 4 h) HSV-2 infection times. For pre-infection ( $-6$  and  $-2$  h), cells were washed thrice by PBS to eliminate excoecarianin prior to the inoculation of the virus. Similar procedures as described in 'antiviral assay' of the 'Methods' section were carried to assess the virus plaques formed.

### Viral Inactivation Assay

The direct effect of excoecarianin on HSV-2 infectivity was evaluated according to previously described procedures (50) with some modifications. Briefly, different concentrations (0.1, 0.5, 1.0, 1.5, 2.0, 2.5 and  $4.0 \mu\text{M}$ ) of excoecarianin were mixed thoroughly with  $1 \times 10^5$  pfu of HSV-2. The mixture was then incubated at  $37^{\circ}\text{C}$  for 6 h. After the incubation, the mixture was diluted at least 100-fold and the residual virus infectivity was determined by plaque assay as described earlier.

### Combined Treatment of ACV and Excoecarianin in Inhibiting HSV-2 Infection

The antiviral activity of excoecarianin in combination with ACV against HSV-2 was evaluated as described previously with some modifications (51–54). The XTT assay was conducted as described above except that HSV-2 at multiplicity of infection (moi) of 1.0 was added concurrently with excoecarianin to the Vero cell monolayer. Inhibitory effect from the combination of ACV and excoecarianin against HSV-2 infection was analyzed by using the isobologram method. The  $\text{IC}_{50}$  was used to calculate the fractional inhibitory concentration (FIC) according to the formula that previously shown (51–54). The interaction between excoecarianin and ACV was interpreted according to the combined FIC index (FIC of excoecarianin plus FIC of ACV). When the combined FIC index is equal to 1, the combination is assumed to act in an additive manner; when it is  $<1$  the interaction is synergistic; and when the combined FIC index is  $>1$  the interaction is antagonistic.

## Statistical Analysis

Data are presented as mean  $\pm$  SD of three independent experiments. The  $IC_{50}$  and  $CC_{50}$  values were calculated by Microsoft Excel 2003. The significance between the test sample and solvent control was analyzed by one-way analysis of variance (ANOVA) followed by Scheffe multiple comparison test. A  $P$  value  $<0.05$  was considered to be statistically significant.

## Results

### Cytotoxicity of Excoecarianin on Vero Cells

The cytotoxic effect of excoecarianin toward Vero cells was investigated by the XTT method. Excoecarianin exhibited cytotoxic effect toward Vero cells in a concentration-dependent manner (Fig. 2) with only 11.57% of cells surviving after treatment at 50  $\mu$ M ( $P < 0.05$ ), but at least 75.2% and 95.4% cells remained alive at concentrations of 5.0 and 1.0  $\mu$ M, respectively. The  $CC_{50}$  of excoecarianin was  $28.0 \pm 2.6 \mu$ M. No significant changes in cellular morphology were observed in excoecarianin-treated cells at a concentration of 3  $\mu$ M and lower (Fig. 3). Overall, the results indicated that excoecarianin showed little cytotoxic effect at concentrations  $<5.0 \mu$ M, and therefore, subsequent studies were performed with excoecarianin treatment at concentrations  $<5.0 \mu$ M.

### Antiviral Activity of Excoecarianin against HSV-1 and HSV-2 Infection


The anti-HSV-1 and HSV-2 activities of excoecarianin were evaluated by plaque reduction assay. Excoecarianin inhibited HSV-2 infection in a concentration-dependent

manner (Fig. 4). The percentage of inhibition were  $0.02 \pm 0.02\%$ ,  $18.3 \pm 8.5\%$ ,  $20.0 \pm 1.5\%$ ,  $56.2 \pm 7.3\%$ ,  $62.8 \pm 11.0\%$ ,  $71.4 \pm 11.9\%$  and  $77.5 \pm 10.8\%$  at the concentrations of 0.1, 0.5, 1.0, 1.5, 2.0, 2.5 and 4.0  $\mu$ M, respectively. The  $IC_{50}$  of excoecarianin against HSV-2 infection was determined to be  $1.4 \pm 0.1 \mu$ M. When tested against HSV-1 infection, excoecarianin was not able to suppress HSV-1 infection (data not shown). At the highest concentration used (4.0  $\mu$ M), excoecarianin could only inhibit 11% of HSV-1 infection.


The selectivity index (SI) measures the safety of a compound to be used as antiviral agent and also confirms that the antiviral effect of a compound is not related to its toxic effect on the cell, is calculated by dividing the  $CC_{50}$  with the  $IC_{50}$  value of the test compound. The SI of excoecarianin was 20.0 against HSV-2 infection.

### Antiviral Activity of Excoecarianin at Different Times of Addition


To study the inhibitory effect of excoecarianin on the stage of HSV-2 infection, the compound was added at different periods (before, during and after) of virus infection. Results showed that excoecarianin added at 2 or 6 h prior to virus infection, and then being removed by PBS washes before the infection, did not have any antiviral activity. When added concurrently with HSV-2, excoecarianin, at concentration of 1.5  $\mu$ M, inhibited 56.2% of virus infection. The inhibitory rate, however, declined to 22.6 and 31.2% when added at 2 and 4 h after infection, respectively (Table 1). This observation indicated that excoecarianin effectively inhibited HSV-2 infection only when it is present at the time of the infection, suggesting its antiviral activity is likely to be at the viral entry stage.



**Figure 2.** The cytotoxic effect of excoecarianin (open bars) and ACV (dotted bars) toward Vero cells as determined by XTT assay. Various concentrations of excoecarianin or ACV were added to Vero cells. After 72h of incubation, the XTT solution was added and then the optical densities were measured. The cytotoxic effect of excoecarianin and ACV were evaluated and the 50% cytotoxic concentration ( $CC_{50}$ ) was calculated. Each bar represents the mean  $\pm$  SD of three independent experiments. The asterisk indicates significant difference between test sample and solvent control ( $P < 0.05$ ).



**Figure 3.** The morphology of Vero cells without treatment (A) and with treatment of ACV (B) or excoecarianin (C and D). Vero cells were seeded onto 24-well culture plates at density of  $1 \times 10^3$  cells per well. After 4 h, excoecarianin or ACV was added. The cells were incubated for 7 days, and the cellular morphology was examined under phase-contrast microscope.



**Figure 4.** Inhibitory effect of excoecarianin (open bars) and ACV (dotted bars) against HSV-2 infection in Vero cell as determined by plaque reduction assay. Vero cells were incubated with 100 pfu of HSV-2 and different concentrations of excoecarianin or ACV. After 1 h, an overlay medium containing 1% methylcellulose was added. On Day 3 post-infection, the cell monolayer was stained with crystal violet and the virus plaques formed were counted. The percentage of inhibition was calculated by comparing the plaque number of compound-treated group to that of the untreated group. The concentrations of excoecarianin and ACV that inhibited 50% of HSV-2 infection ( $\text{IC}_{50}$ ) were determined. Each bar represents the mean  $\pm$  SD of three independent experiments. The asterisk indicates significant difference between test sample and solvent control ( $P < 0.05$ ).

#### Mechanism of Action of Excoecarianin against HSV-2 Infection

To investigate whether excoecarianin inhibited HSV-2 through inactivation of viral particles, a suspension of

the virus was treated at  $37^\circ\text{C}$  for 6 h with various concentrations of excoecarianin. The residual viral infectivity was determined by plaque assay. As shown in Table 2, excoecarianin inactivated viral infectivity in

**Table 1.** The effect of treatment time on the antiviral activity of excoecarianin

Time periods of compound addition	Percentage of inhibition	
	Excoecarianin	ACV
Before infection		
−6	0.0 ± 0.0	0.0 ± 0.0
−2	0.0 ± 0.0	15.0 ± 7.6
During infection		
0	56.2 ± 7.3*	95.4 ± 1.7*
After infection		
2	22.6 ± 10.2	92.6 ± 5.0*
4	31.2 ± 19.2	94.9 ± 4.7*

Excoecarianin and ACV were tested at a concentration of 1.5 and 5  $\mu\text{M}$ , respectively; Data were mean  $\pm$  SD of three independent experiments; \* $P < 0.05$  (compared between control and tested compound).

**Table 2.** The effect of excoecarianin against HSV-2 infectivity

	Concentration ( $\mu\text{M}$ )	Percentage control of viral infectivity
Control	0	100.0 $\pm$ 0.0
Excoecarianin	0.1	91.6 $\pm$ 7.4
	0.5	98.8 $\pm$ 2.1
	1.0	96.2 $\pm$ 6.6
	1.5	61.3 $\pm$ 7.8*
	2.0	27.7 $\pm$ 7.9*
	2.5	36.3 $\pm$ 10.8*
	4.0	1.5 $\pm$ 0.1*
ACV	0.5	100.0 $\pm$ 0.5
	2.5	92.8 $\pm$ 6.5
	5.0	92.9 $\pm$ 4.9
	10.0	91.7 $\pm$ 8.2

Excoecarianin was mixed with HSV-2 at 37°C for 6 h before the residual viral titer was determined by plaque assay; Values were mean  $\pm$  SD of three independent experiments; \* $P < 0.05$  (compared between control and tested compound).

concentration-dependent manner. While it did not inactivate HSV-2 at concentration of  $\leq 1.0 \mu\text{M}$ , the residual viral infectivity gradually declined from 61.3  $\pm$  7.8%, to 27.7  $\pm$  7.9%, and to 36.3  $\pm$  10.8% as the concentrations of excoecarianin increased from 1.5, to 2.0, and to 2.5  $\mu\text{M}$ , respectively. Excoecarianin drastically reduced viral infectivity at concentration of 4.0  $\mu\text{M}$  with only 1.5  $\pm$  0.1% of virus being able to produce infectivity ( $P < 0.05$ ). In contrast, ACV, a well-defined anti-HSV-2 agent which specifically inhibits viral DNA synthesis, did not have effect on virus infectivity even at a concentration of 10.0  $\mu\text{M}$ .


### Combined Treatment of ACV and Excoecarianin against HSV-2 Infection

The inhibitory effect of ACV plus excoecarianin against HSV-2 infection in Vero cells was examined by XTT assay. The inhibitory activity was further evaluated by the isobologram method. Low concentrations of ACV

**Table 3.** Inhibitory effects of excoecarianin in combination with ACV against the infection of HSV-2 in Vero cells

Compound(s)	Mean ( $\text{IC}_{50} \pm \text{SD}$ ) <sup>a</sup>	$\text{FIC}_{\text{excoecarianin}} + \text{FIC}_{\text{ACV}}$ <sup>b</sup>	FIC index interpretation
Excoecarianin alone	11.87 $\pm$ 2.9	—	—
ACV alone	1.33 $\pm$ 0.84	—	—
ACV + 1.0 $\mu\text{M}$ excoecarianin	0.76 $\pm$ 0.34	0.66	synergistic
ACV + 2.5 $\mu\text{M}$ excoecarianin	0.39 $\pm$ 0.18	0.51	synergistic
ACV + 5.0 $\mu\text{M}$ excoecarianin	0.11 $\pm$ 0.02	0.50	synergistic

<sup>a</sup>Results are based on three independent experiments; <sup>b</sup> $\text{FIC}_{\text{excoecarianin}}$  and  $\text{FIC}_{\text{ACV}}$  are the fractional inhibitory concentration (FIC) of excoecarianin and ACV, respectively.



**Figure 5.** Synergistic antiviral activity of ACV and excoecarianin in Vero cells.  $\text{IC}_{50}$  values were derived from the data shown in Table 3 and used to construct the isobologram.  $\text{FIC}_{50}$  of ACV represents the ratio of the  $\text{IC}_{50}$  of ACV in the presence of a constant concentration of excoecarianin to the  $\text{IC}_{50}$  of ACV alone. The X-axis represents the ratio of the fixed concentration of excoecarianin to the  $\text{IC}_{50}$  of excoecarianin alone. In this representation, displacement of the experimental data points to the left of the theoretical line is indicative of synergistic behavior.

could inhibit HSV-2 infection by the addition of excoecarianin (Table 3). The  $\text{IC}_{50}$  value for ACV could be reduced from 1.33  $\pm$  0.84  $\mu\text{M}$  of ACV alone to 0.76  $\pm$  0.34, 0.39  $\pm$  0.18 and 0.11  $\pm$  0.02  $\mu\text{M}$  of ACV plus 1.0, 2.5 and 4.0  $\mu\text{M}$  of excoecarianin, respectively. The FICs of ACV plus excoecarianin were in the range 0.50–0.66, indicating that the combinatorial effect of ACV plus excoecarianin in inhibiting HSV-2 infection was synergistic (Table 3 and Fig. 5). None of these drug combinations exhibited cytotoxic effect against Vero cells at the concentrations used as assessed by cytotoxicity assay (data not shown).


## Discussion

Our present study demonstrated that excoecarianin inhibited HSV-2 infection but not HSV-1 infection. This antiviral effect was not correlated to cellular cytotoxicity as the effective micromolar concentrations of excoecarianin used did not affect cell viability (Fig. 2) or induce changes in cellular morphology (Fig. 3). The antiviral activity was most prominent when excoecarianin was added during viral infection. This effect was attributed to the compound's ability to directly reduce viral infectivity through interaction with HSV-2 viral particles. In addition, combined treatment of ACV and excoecarianin synergistically inhibited HSV-2 infection.

Our time-point assessment of antiviral activity indicated that excoecarianin most effectively inhibited HSV-2 infection only when it was concurrently present with the virus at the time of infection, but was less effective when cells were pre-treated with it (and washed) or treated post viral entry. Since our pre-treatment protocol involved washing the compound off prior to infection, these results would speculate that excoecarianin probably does not prevent HSV-2 infection by masking cell surface HSV-2 receptors through covalent bonds formation to remain attached, nor does it trigger any cellular antiviral responses including induction of type I interferons to control the viral infection. Clearly, excoecarianin is affecting the early stage of HSV-2 infection and the

presence of the drug at the time of the infection is crucial. The observation suggests that the inhibition on HSV-2 by excoecarianin was most likely mediated by interfering with virus entry step without having any direct effect on the cell itself, and indeed it was confirmed by our subsequent data that excoecarianin directly inactivated HSV-2 particles (Table 2). This effect is different from that of ACV, which acts as an inhibitor of HSV DNA synthesis. The ability of excoecarianin to inactivate HSV-2 virus particles thus makes it useful as an anti-HSV-2 agent in preventing *de novo* viral infection and thereby could help control viral spread and limit recurrent infections.

In virus taxonomy, HSV-1 and HSV-2 are members of the genera *Simplexvirus* from the subfamily *Alphaherpesvirinae* in *Herpesviridae* family (55). Both viruses are sensitive to nucleoside-like drugs, such as ACV. The effectiveness on inhibiting HSV-2 but not HSV-1 infection suggested that excoecarianin is specific in mediating its anti-HSV activity. As HSV-1 and HSV-2 virions share high degree of similarity, it is unclear at the moment how excoecarianin is specific only to HSV-2. Excoecarianin could possibly be specific to viral glycoproteins expressed by HSV-2 such as glycoprotein C (gC-2) which has been shown to contribute to serotype differences in cell tropism and may play a different role with respect to its HSV-1 counterpart gC-1 (56,57).



**Figure 6.** The time-of-addition effect of excoecarianin against HSV-2 infection. Excoecarianin added prior to virus infection and then washed out did not protect the cells from HSV-2 infection. The addition of excoecarianin after 2 h of the virus inoculation also did not inhibit HSV-2 infection. The compound can inhibit HSV-2 infection only when it was added concurrently with the virus infection.

Further studies are required to clarify the underlying reason(s) of the selectivity in anti-HSV effects mediated by excoecarianin.

Our previous studies also revealed that the acetone extract and several pure compounds found in the extracts from *P. urinaria* could inhibit HSV-2 infection at different magnitudes of potency (44–46). Particularly, we have observed that the acetone extract of *P. urinaria* inhibited HSV-2 infection through diminishing the virus infectivity, and the acetone extract was only active in inhibiting HSV-2 when added concurrently with the virus (44). The acetone extract also did not have any direct effect on the cells in preventing viral infection as indicated by a lack of benefit from pre-treatment dosage (44). Based on our current experimental observations, the acetone-isolated excoecarianin from *P. urinaria* is likely responsible, at least in part, for the previous antiviral effects observed from the *P. urinaria* acetone extract. Furthermore, the identification of excoecarianin as a specific HSV-2 entry inhibitor by our study suggests that excoecarianin or the standardized acetone extract *P. urinaria* could both be explored for therapeutic development against HSV-2 infection. As *P. urinaria* also possesses antiviral activity against other viruses as well, including hepatitis B virus (37–41), retrovirus (42) and Epstein-Barr virus (43), it is unclear whether excoecarianin can mediate similar antiviral activities against other viruses. It would also be interesting to investigate whether excoecarianin or the standardized extract could improve the disease pathogenesis in viral co-infection cases such as in immunocompromised patients as an adjunct therapy.

We have demonstrated here that the combined treatment of ACV and excoecarianin exhibits synergistic activity in inhibiting HSV-2 infection, decreasing the concentration required for ACV to attain similar antiviral profile. This result has clinical significance in the management of HSV-2 infection. As there are more ACV-resistant HSV related cases being identified from immunocompromised patients as a result of the emergence of HIV or organ transplantation (6,7,58), anti-HSV agents targeting viral enzymes or factors essential for entry will likely be useful for controlling nucleoside-resistant strains (59). The application of excoecarianin would complement ACV treatment, not only in lowering the amount of drug necessary to achieve viral eradication but would also help decrease the risk of ACV-resistance from occurring. Thus, the difference in mechanism of action between excoecarianin and ACV, along with the synergistic effect observed from their combined treatment, implies that excoecarianin could be useful in therapy against HSV-2 infection in immunocompromised patients.

In summary, we have identified the acetone-extracted excoecarianin from *P. urinaria* as an antiviral compound that is specific for HSV-2, which may be responsible in part for the previously observed *P. urinaria*'s

antiviral effect against this viral infection. The ability of excoecarianin to inactivate HSV-2 virus particles and to act synergistically with the nucleoside analogue ACV may make it useful to control viral spread in *de novo* and recurrent infections. We therefore suggest that the development of excoecarianin or the standardized extract of *P. urinaria* as an oral agent or as a topical cream may be further investigated as an effective strategy in managing HSV-2 infections.

## Funding

National Science Council grant [NSC 92-2320-B-037-044].

## References

- Kleymann G, Fischer R, Betz UA, Hendrix M, Bender W, Schneider U, et al. New helicase-primase inhibitors as drug candidates for the treatment of herpes simplex disease. *Nat Med* 2002;8:392–98.
- Liuzzi M, Deziel R, Moss N, Beaulieu P, Bonneau AM, Bousquet C, et al. A potent peptidomimetic inhibitor of HSV ribonucleotide reductase with antiviral activity *in vivo*. *Nature* 1994;372:695–98.
- Wathen MW. Non-nucleoside inhibitors of herpesviruses. *Rev Med Virol* 2002;12:167–78.
- De Clercq E. Molecular targets for antiviral agents. *J Pharmacol Exp Ther* 2001;297:1–10.
- Naesens L, De Clercq E. Recent developments in herpesvirus therapy. *Herpes* 2001;8:12–16.
- Stranska R, Schuurman R, Nienhuis F, Goedegebuure IW, Polman M, Weel JF, et al. Survey of acyclovir-resistant herpes simplex virus in the Netherlands: prevalence and characterization. *J Clin Virol* 2005;32:7–18.
- Ziyaeyan M, Alborzi A, Japoni A, Kadivar M, Davarpanah MA, Pourabbas B, et al. Frequency of acyclovir-resistant herpes simplex viruses isolated from the general immunocompetent population and patients with acquired immunodeficiency syndrome. *Int J Dermatol* 2007;46:1263–66.
- Frangoul H, Wills M, Crossno C, Engel M, Domm J. Acyclovir-resistant herpes simplex virus pneumonia post-unrelated stem cell transplantation: a word of caution. *Pediatr Transplant* 2007;11:942–44.
- Griffiths PD. Herpesviruses and AIDS. *Scand J Infect Dis* 1996;Suppl 100:3–7.
- Pillay D, Mutimer D, Singhal S, Turner A, Ward K, Wood M. Management of herpes virus infections following transplantation. *J Antimicrob Chemother* 2000;45:729–48.
- Cooper EL. Drug discovery, CAM and natural products. *eCAM* 2001;1:215–17.
- Mukhtar M, Arshad M, Ahmad M, Pomerantz RJ, Wigdahl B, Parveen Z. Antiviral potentials of medicinal plants. *Virus Res* 2008;131:111–20.
- Rajbhandari M, Mentel R, Jha PK, Chaudhary RP, Bhattarai S, Gewali MB, et al. Antiviral activity of some plants used in Nepalese traditional medicine. *eCAM* doi:10.1093/ecam/nem156.
- Saddi M, Sanna A, Cottiglia F, Chisu L, Casu L, Bonsignore L, et al. Antitherpesvirus activity of *Artemisia arborescens* essential oil and inhibition of lateral diffusion in Vero cells. *Ann Clin Microbiol Antimicrob* 2007;6:10.
- Chularojmontri L, Wattanapitayakul SK, Herunsalee A, Charuchongkolwongse S, Niumsukul S, Srichairat S. Antioxidative and cardioprotective effects of *Phyllanthus urinaria* L. on doxorubicin-induced cardiotoxicity. *Biol Pharm Bull* 2005;28:1165–71.
- Dias MA, Campos AH, Cechinel-Filho V, Yunes RA, Calixto JB. Analysis of the mechanisms underlying the contractile responses induced by the hydroalcoholic extract of *Phyllanthus urinaria* in



- the guinea pig urinary bladder *in-vitro*. *J Pharm Pharmacol* 1995;47:846–851.
17. Fang SH, Rao YK, Tzeng YM. Anti-oxidant and inflammatory mediator's growth inhibitory effects of compounds isolated from *Phyllanthus urinaria*. *J Ethnopharmacol* 2008;116:333–40.
  18. Giridharan P, Somasundaram ST, Perumal K, Vishwakarma RA, Karthikeyan NP, Velmurugan R, et al. Novel substituted methylenedioxy lignan suppresses proliferation of cancer cells by inhibiting telomerase and activation of c-myc and caspases leading to apoptosis. *Br J Cancer* 2002;87:98–105.
  19. Higashino H, Suzuki A, Tanaka Y, Pootakham K. Hypoglycemic effects of Siamese *Momordica charantia* and *Phyllanthus urinaria* extracts in streptozotocin-induced diabetic rats. *Nippon Yakurigaku Zasshi* 1992;100:415–21.
  20. Huang ST, Yang RC, Yang LJ, Lee PN, Pang JH. *Phyllanthus urinaria* triggers the apoptosis and Bcl-2 down-regulation in Lewis lung carcinoma cells. *Life Sci* 2003;72:1705–16.
  21. Huang ST, Yang RC, Chen MY, Pang JH. *Phyllanthus urinaria* induces the Fas receptor/ligand expression and ceramide-mediated apoptosis in HL-60 cells. *Life Sci* 2004;75:339–51.
  22. Huang ST, Yang RC, Pang JH. Aqueous extract of *Phyllanthus urinaria* induces apoptosis in human cancer cells. *Am J Chin Med* 2004;32:175–83.
  23. Huang ST, Yang RC, Lee PN, Yang SH, Liao SK, Chen TY. Anti-tumor and anti-angiogenic effects of *Phyllanthus urinaria* in mice bearing Lewis lung carcinoma. *Int Immunopharmacol* 2006;6:870–9.
  24. Lai CH, Fang SH, Rao YK, Geethangili M, Tang CH, Lin YJ, et al. Inhibition of *Helicobacter pylori*-induced inflammation in human gastric epithelial AGS cells by *Phyllanthus urinaria* extracts. *J Ethnopharmacol* 2008;118:522–6.
  25. Lans CA. Ethnomedicines used in Trinidad and Tobago for urinary problems and diabetes mellitus. *J Ethnobiol Ethnomed* 2006;2:45.
  26. Lans CA. Comparison of plants used for skin and stomach problems in Trinidad and Tobago with Asian ethnomedicine. *J Ethnobiol Ethnomed* 2007;3:3.
  27. Lee CY, Peng WH, Cheng HY, Chen FN, Lai MT, Chiu TH. Hepatoprotective effect of *Phyllanthus* in Taiwan on acute liver damage induced by carbon tetrachloride. *Am J Chin Med* 2006;34:471–82.
  28. Lin SY, Wang CC, Lu YL, Wu WC, Hou WC. Antioxidant, anti-semicarbazide-sensitive amine oxidase, and anti-hypertensive activities of geraniin isolated from *Phyllanthus urinaria*. *Food Chem Toxicol* 2008;46:2485–92.
  29. Paulino N, Cechinel-Filho V, Pizzolatti MG, Yunes RA, Calixto JB. Mechanisms involved in the contractile responses induced by the hydroalcoholic extract of *Phyllanthus urinaria* on the guinea pig isolated trachea: evidence for participation of tachykinins and influx of extracellular Ca<sup>2+</sup> sensitive to ruthenium red. *Gen Pharmacol* 1996;27:795–802.
  30. Paulino N, Cechinel-Filho V, Yunes RA, Calixto JB. The relaxant effect of extract of *Phyllanthus urinaria* in the guinea-pig isolated trachea. Evidence for involvement of ATP-sensitive potassium channels. *J Pharm Pharmacol* 1996;48:1158–63.
  31. Santos AR, Cechinel-Filho V, Yunes RA, Calixto JB. Analysis of the mechanisms underlying the antinociceptive effect of the extracts of plants from the genus *Phyllanthus*. *Gen Pharmacol* 1995;26:1499–506.
  32. Santos AR, De Campos RO, Miguel OG, Cechinel-Filho V, Yunes RA, Calixto JB. The involvement of K<sup>+</sup> channels and Gi/o protein in the antinociceptive action of the gallic acid ethyl ester. *Eur J Pharmacol* 1999;379:7–17.
  33. Shen B, Yu J, Wang S, Chu ES, Wong VW, Zhou X, et al. *Phyllanthus urinaria* ameliorates the severity of nutritional steatohepatitis both *in vitro* and *in vivo*. *Hepatology* 2008;47:473–83.
  34. Xu M, Zha ZJ, Qin XL, Zhang XL, Yang CR, Zhang YJ. Phenolic antioxidants from the whole plant of *Phyllanthus urinaria*. *Chem Biodivers* 2007;4:2246–52.
  35. Zhang DY, He DW, Wei GH, Liu X, Lin T, Li XL. *Phyllanthus urinaria* extract promotes N-cadherin expression in nitrogen mustard-disrupted testicular tissues *in vivo*. *Zhonghua Nan Ke Xue* 2008;14:396–400 (in Chinese).
  36. Zhou S, Xu C, Zhou N, Huang Y, Huang L, Chen X, et al. Mechanism of protective action of *Phyllanthus urinaria* L. against injuries of liver cells. *Zhongguo Zhong Yao Za Zhi* 1997;22:109–11 (in Chinese).
  37. Chen YX, Guo SH, Zhang DF. Experimental study on anti-duck hepatitis B viral effect of *Phyllanthus urinaria* of different areas and combined therapy with other drugs. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 1995;15:225–7 (in Chinese).
  38. Ji XH, Qin YZ, Wang WY, Zhu JY, Liu XT. Effects of extracts from *Phyllanthus urinaria* L. on HbsAg production in PLC/PRF/5 cell line. *Zhongguo Zhong Yao Za Zhi* 1993;18:496–98, 511 (in Chinese).
  39. Wang M, Cheng H, Li Y, Meng L, Zhao G, Mai K. Herbs of the genus *Phyllanthus* in the treatment of chronic hepatitis B: observations with three preparations from different geographic sites. *J Lab Clin Med* 1995;126:350–52.
  40. Wang MX, Cheng HW, Li YJ, Meng LM, Mai K. Efficacy of *Phyllanthus* spp. in treating patients with chronic hepatitis B. *Zhongguo Zhong Yao Za Zhi* 1994;19:750–1 764 (in Chinese).
  41. Zhong Y, Zuo C, Li F, Ding X, Yao Q, Wu K, et al. Chemical constituents of *Phyllanthus urinaria* L. and its antiviral activity against hepatitis B virus. *Zhongguo Zhong Yao Za Zhi* 1998;23:363–4. 384 (in Chinese).
  42. Suthienkul O, Miyazaki O, Chulasiri M, Kositanont U, Oishi K. Retroviral reverse transcriptase inhibitory activity in Thai herbs and spices: screening with Moloney murine leukemia viral enzyme. *Southeast Asian J Trop Med Public Health* 1993;24:751–5.
  43. Liu KC, Lin MT, Lee SS, Chiou JF, Ren S, Lien EJ. Antiviral tannins from two *Phyllanthus* species. *Planta Med* 1999;65:43–6.
  44. Yang CM, Cheng HY, Lin TC, Chiang LC, Lin CC. Acetone, ethanol and methanol extracts of *Phyllanthus urinaria* inhibit HSV-2 infection *in vitro*. *Antiviral Res* 2005;67:24–30.
  45. Yang CM, Cheng HY, Lin TC, Chiang LC, Lin CC. The *in vitro* activity of geraniin and 1, 3, 4, 6-tetra-O-galloyl- $\beta$ -D-glucose isolated from *Phyllanthus urinaria* against herpes simplex virus type 1 and type 2 infection. *J Ethnopharmacol* 2007;110:555–8.
  46. Yang CM, Cheng HY, Lin TC, Chiang LC, Lin CC. Hippomanin A from acetone extract of *Phyllanthus urinaria* inhibited HSV-2 but not HSV-1 infection *in vitro*. *Phytother Res* 2007;21:1182–6.
  47. Lin JH, Tanaka T, Nonaka GI, Nishioka I, Chen IS. Tannins and related compounds. XCVIII.: Structure of three new dimeric ellagitannins, excoecarianin and excoecarinins A and B, isolated from the leaves of *Excoecaria kawakamii* HAYATA. *Chem Pharm Bull* 1990;38:2162–71.
  48. Burleson FG, Chambers TM, Wiedbrauk DL. Plaque assay. In: Burleson FG, Chambers TM, Wiedbrauk DL (eds). *Virology: A Laboratory Manual*. California: Academic Press Inc, 1992, 74–84.
  49. Cheng HY, Lin LT, Huang HH, Yang CM, Lin CC. Yin Chen Hao Tang, a Chinese prescription, inhibits both herpes simplex virus type-1 and type-2 infection *in vitro*. *Antiviral Res* 2008;77:14–9.
  50. Cheng HY, Yang CM, Lin TC, Shieh D E, Lin CC. ent-Epiatzelechin-(4 $\alpha$ →8)-epiatzelechin extracted from *Cassia javanica* inhibits herpes simplex virus type 2 replication. *J Med Microbiol* 2006;55:201–6.
  51. Cheng HY, Lin TC, Yang CM, Wang KC, Lin LT, Lin CC. Putranjivain A from *Euphorbia jolkini* inhibits both virus entry and late stage replication of herpes simplex virus type 2 *in vitro*. *J Antimicrob Chemother* 2004;53:577–83.
  52. Lawetz C, Liuzzi M. The antiviral activity of the ribonucleotide reductase inhibitor BILD 1351 SE in combination with acyclovir against HSV type-1 in cell culture. *Antiviral Res* 1998;39:35–46.
  53. Piret J, Roy S, Gagnon M, Landry S, Desormeaux A, Omar RF, et al. Comparative study of mechanism of herpes simplex virus inactivation by sodium lauryl sulfate and n-lauroylsarcosine. *Antimicrob Agents Chemother* 2002;46:2933–42.
  54. Suhnel J. Evaluation of synergism or antagonism for the combined action of antiviral agents. *Antiviral Res* 1990;13:23–39.
  55. Roizman B, Pellet PE. The family Herpesviridae: a brief introduction. In: Knipe DM, Howley PM, Griffin DE, Martin MA, Lamb RA, Roizman B, et al. (eds). *Fields' Virology*. Philadelphia: Lippincott Williams and Wilkins, 2001, 2381–97.
  56. Gerber SI, Belval BJ, Herold BC. Differences in the role of glycoprotein C of HSV-1 and HSV-2 in viral binding may contribute to serotype differences in cell tropism. *Virology* 1995;214:29–39.

57. Rux AH, Lou H, Lambris JD, Friedman HM, Eisenberg RJ, Cohen GH. Kinetic analysis of glycoprotein C of herpes simplex virus types 1 and 2 binding to heparin, heparan sulfate, and complement component C3b. *Virology* 2002;294:324–32.
58. Danve-Szatanek C, Aymard M, Thouvenot D, Morfin F, Agius G, Bertin I, et al. Surveillance network for herpes simplex virus resistance to antiviral drugs: 3-year follow-up. *J Clin Microbiol* 2004;42:242–9.
59. Cassady KA, Whitley RJ. New therapeutic approaches to the alpha-herpesvirus infections. *J Antimicrob Chemother* 1997;39:119–28.

Received March 26, 2009; accepted August 24, 2009